

Stability of line bundles transforms on curves with respect to low codimensional subspaces

Ernesto Mistretta

Abstract

We show the stability of certain syzygies of line bundles on curves, which we call transforms, and are kernels of the evaluation map on subspaces of the space of global sections. For the transforms constructed, we prove the existence of reducible theta divisors, in the cases where the slope is integer.

1 Introduction

In the study of vector bundles on curves, it is a natural question to investigate the stability of kernels of evaluation maps of global sections. This was used in particular by Paranjape and Ramanan (cf. [PR88]), and Butler (cf. [But94]), to prove normal generation of certain vector bundles, by Ein and Lazarsfeld (cf. [EL92]) to show the stability of the Picard bundle, and by Beauville (*e.g.* in [Bea03]) to study theta divisors.

Definition 1.1 *Let C be a smooth projective curve over an algebraically closed field $\mathbb{k}$, and E a globally generated vector bundle over C . We call $M_{V,E} := \ker(V \otimes \mathcal{O}_C \rightarrow E)$ the transform of the vector bundle E with respect to the generating subspace $V \subset H^0(C, E)$, and $M_E := M_{H^0(C, E), E} = \ker(H^0(C, E) \otimes \mathcal{O}_C \rightarrow E)$ the total transform of E .*

Starting from a result of Butler, who proved the stability of total transforms under certain hypothesis, we want to investigate the stability of transforms of line bundles by generic subspaces of certain codimensions.

Theorem 1.2 (Butler) *Let C be a smooth projective curve of genus $g \geq 1$ over an algebraically closed field $\mathbb{k}$, and E a semistable vector bundle over C with slope $\mu(E) \geq 2g$, then the vector bundle $M_E := \ker(H^0(C, E) \otimes \mathcal{O}_C \rightarrow E)$ is semistable. Furthermore, if E is stable and $\mu(E) \geq 2g$, then M_E is stable, unless $\mu(E) = 2g$, and either C is hyperelliptic or $\omega_C \hookrightarrow E$.*

It is natural to ask what happens taking subspaces in the place of the vector space of global sections. Our results can be resumed to the following theorem:

Theorem 1.3 *Let $\mathcal{L}$ be a line bundle of degree d on a curve C of genus $g \geq 2$, such that $d \geq 2g + 2c$, with $1 \leq c \leq g$. Then $M_{V,\mathcal{L}}$ is semistable for a generic subspace $V \subset H^0(\mathcal{L})$ of codimension c . It is stable unless $d = 2g + 2c$ and the curve is hyperelliptic.*

Similar results can be deduced by some constructions in Vincent Mercat's work [Mer99] on Brill-Noether's loci, but we think that in our case it is useful to give a more direct proof which applies to all line bundles of degree $d \geq 2g + 2c$ and not only generic ones.

Eventually, we observe the existence of theta divisors associated to the (semi)stable transforms having integer slope -2 . Those theta divisors are always non integral, and in most cases reducible, hence give further examples of stable vector bundles admitting a reducible theta divisor (cf. [Bea03]).

Remark 1.4 A geometrical interpretation of those kinds of results goes as follows: a generating subspace $V \subset H^0(C, \mathcal{L})$ gives rise to a base point free linear system $|V| \subset |\mathcal{L}|$ on the curve C , and determines a map $\varphi_V: C \rightarrow \mathbb{P}(V^*)$, which associates to a point $x \in C$ the hyperplane of global sections in V vanishing in x . The Euler sequence on $\mathbb{P}(V^*)$ is the dual of the tautological sequence:

$$0 \rightarrow \Omega_{\mathbb{P}(V^*)}(1) \rightarrow V \otimes \mathcal{O}_{\mathbb{P}(V^*)} \rightarrow \mathcal{O}_{\mathbb{P}(V^*)}(1) \rightarrow 0$$

which restricted to C gives the evaluation sequence

$$0 \rightarrow M_{V,\mathcal{L}} \rightarrow V \otimes \mathcal{O}_C \rightarrow \mathcal{L} \rightarrow 0.$$

As stability of a vector bundle is not affected by dualizing and tensorizing by a line bundle, we see that stability of $M_{V,\mathcal{L}} = \Omega(1)|_C$ is equivalent to the stability of the restriction of the tangent bundle of the projective space $\mathbb{P}(V^*)$ to the curve C .

So our theorem translates to

Theorem 1.5 *Let $C \subset \mathbb{P}^{d-g}$ be a genus $g \geq 2$ degree d non-degenerate smooth curve, where $d > 2g + 2c$, and c is a constant such that $1 \leq c \leq g$. Then for the generic projection $\mathbb{P}^{d-g} \dashrightarrow \mathbb{P}^{d-g-c}$ the restriction $T_{\mathbb{P}^{d-g-c}|_C}$ is stable.*

1.1 Acknowledgments

I wish to thank my advisor Daniel Huybrechts for his help during my thesis, Arnaud Beauville for his interest and his comments, and my colleagues of Paris and Rome for their friendly advices.

2 Stability of transforms

We essentially use the following two lemmas:

Lemma 2.1 (Butler) *Let C be a curve of genus $g \geq 2$, F a vector bundle on C with no trivial summands, and such that $h^1(F) \neq 0$. Suppose that $V \subset H^0(F)$ generates F . If $N = M_{V,F}$ is stable, then $\mu(N) \leq -2$. Furthermore, $\mu(N) = -2$ implies that either C is hyperelliptic F is the hyperelliptic bundle and N its dual, or $F = \omega$ and $N = M_\omega$.*

The proof of this lemma is based on the result by Paranjape, Ramanan asserting the stability of M_ω (see [But94] and [PR88]).

Lemma 2.2 *Let $\mathcal{L}$ be a degree $d \geq 2g + 2c$ line bundle on a curve C of genus $g \geq 2$, with $c \leq g$ and let $V \subset H^0(\mathcal{L})$ be a generating subspace of codimension c . Suppose there exists a stable subbundle of maximal slope $N \hookrightarrow M_{V,\mathcal{L}}$ such that $0 \neq N \neq M_{V,\mathcal{L}}$ and $\mu(N) \geq \mu(M_{V,\mathcal{L}})$.*

Then there exists a line bundle F of degree $f \leq d - 1$, a generating subspace $W \subset H^0(F)$, and an injection $F \hookrightarrow \mathcal{L}$ such that N fits into the following commutative diagram

$$\begin{array}{ccccccc} 0 & \rightarrow & N & \rightarrow & W \otimes \mathcal{O}_C & \rightarrow & F \rightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \rightarrow & M_{V,\mathcal{L}} & \rightarrow & V \otimes \mathcal{O}_C & \rightarrow & \mathcal{L} \rightarrow 0, \end{array}$$

i.e. a destabilization of $M_{V,\mathcal{L}}$ must be the transform of a line bundle injecting into $\mathcal{L}$ such that the global sections we are transforming by are in V .

The importance of this lemma lies in the fact that we associate a line bundle F to a destabilizing N , and this allows us more easily to parametrize destabilizations and bound their dimension.

Proof We remark that $\mu(M_{V,\mathcal{L}}) = -d/(d - g - c) \geq -2$ for $d \geq 2g + 2c$. Consider a stable subbundle $N \hookrightarrow M_{V,\mathcal{L}}$ of maximal slope. Then it fits into the commutative diagram

$$\begin{array}{ccccccc} 0 & \rightarrow & N & \rightarrow & W \otimes \mathcal{O}_C & \rightarrow & F \rightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \rightarrow & M_{V,\mathcal{L}} & \rightarrow & V \otimes \mathcal{O}_C & \rightarrow & \mathcal{L} \rightarrow 0 \end{array}$$

where $W \hookrightarrow V$ is defined by $W^* := \text{Im}(V^* \rightarrow H^0(N^*))$, hence W^* generates N^* , and we call $F^* := \ker(W^* \otimes \mathcal{O} \rightarrow N^*)$.

Then F is a vector bundle with no trivial summands. Moreover the morphism $F \rightarrow \mathcal{L}$ is not zero, as $W \otimes \mathcal{O}$ does not map to $M_{V,\mathcal{L}}$. We have to show that $\text{rk} F = 1$ and that $W = H^0(F)$. We distinguish the two cases $h^1(F) = 0$ or $h^1(F) \neq 0$.

• Let us suppose that $h^1(F) = 0$. Then $h^0(F) = \chi(F) = \text{rk} F(\mu(F) + 1 - g)$. On the other hand, $h^0(F) > \text{rk} F$ as F is globally generated and not trivial.

Together this yields

$$\mu(F) > g. \quad (1)$$

Furthermore

$$\mu(N) = -\deg F / (\dim W - \text{rk} F) \leq -\mu(F) / (\mu(F) - g) = \mu(M_F), \quad (2)$$

as $\dim W \leq h^0(F) = \operatorname{rk} F(\mu(F) + 1 - g)$.

Consider the image $I = \operatorname{Im}(F \rightarrow \mathcal{L}) \subseteq \mathcal{L}$. The commutative diagram

$$\begin{array}{ccccc} W & \hookrightarrow & H^0(F) & \rightarrow & H^0(I) \\ \downarrow & & & & \downarrow \\ V & & \hookrightarrow & & H^0(\mathcal{L}) \end{array}$$

shows that the map $W \rightarrow H^0(I)$ is injective and its image $W' \subset H^0(I)$ is contained in $V \subset H^0(\mathcal{L})$, hence $N \hookrightarrow M_{W',I} \hookrightarrow M_{V,\mathcal{L}}$. As N is a subbundle of $M_{V,\mathcal{L}}$ of maximal slope, this yields $\mu(N) \geq \mu(M_{W',I})$, *i.e.* $-\deg F/\operatorname{rk} N \geq -\deg I/\operatorname{rk} M_{W',I}$. Then

$$\deg F \leq \deg I(\operatorname{rk} N/\operatorname{rk} M_{W',I}) \leq \deg I \leq \deg \mathcal{L} = d .$$

If $\operatorname{rk} F \geq 2$, then $\mu(F) \leq \deg \mathcal{L}/2 = d/2$, so

$$\mu(N) \leq \frac{-\mu(F)}{\mu(F) - g} \leq \frac{-d/2}{d/2 - g} = \frac{-d}{d - 2g} \leq \frac{-d}{d - g - c} = \mu(M_{V,\mathcal{L}}) .$$

Here the first inequality is (2). For the second one shows that the function $-x/(x - g)$ is strictly increasing for $x > g$. Then use $\mu(F) > g$ due to (1). Equality holds only if $\operatorname{rk} F = 2$, $\deg F = d$, $W = H^0(F)$, and $g = c$. But in this case we would find that $\dim W = h^0(F) = d + 2 - 2g > d + 1 - g - c = \dim V$, which is impossible as by construction $W \hookrightarrow V$.

Hence $\operatorname{rk} F = 1$. So $F = I$ is a globally generated and acyclic line bundle of degree $f \leq d$, and $\mu(N) = -f/(\dim W - 1)$.

It is easy to see that the case $f = d$ cannot hold, as in that case we cannot have $\mu(N) \geq \mu(M_{V,\mathcal{L}})$. So $f \leq d - 1$.

- In the case $h^1(F) \neq 0$, by lemma 2.1, $\mu(N) \leq -2$. Equality holds only if $F = \omega_C$ and $W = H^0(\omega)$, or if the curve C is hyperelliptic and F is the hyperelliptic bundle. In the latter case the only generating space of global sections is $H^0(F)$. In any case we have $f = \deg F < d - 1$. $\square$

Remark 2.3 The diagram in the statement of the lemma is a construction from Butler's proof of theorem 1.2.

Remark 2.4 Looking carefully at the numerical invariants in the above proof, we can deduce some inequalities which will be useful in the following: let us consider again the diagram in the above lemma

$$\begin{array}{ccccccc} 0 & \rightarrow & N & \rightarrow & W \otimes \mathcal{O}_C & \rightarrow & F \rightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \rightarrow & M_{V,\mathcal{L}} & \rightarrow & V \otimes \mathcal{O}_C & \rightarrow & \mathcal{L} \rightarrow 0 \end{array}$$

and suppose that $h^1(F) = 0$. Let us call $f := \deg F$, $s := d - f$, and $b := \operatorname{codim}_{H^0(F)} W$. Then we can show that

$$0 < c - b < s \leq \frac{d}{g + c}(c - b) . \quad (3)$$

In fact, as $W \hookrightarrow V$, and $W \neq V$, then

$$d - s + 1 - g - b = h^0(F) - b = \dim W < \dim V = d + 1 - g - c ,$$

hence $c - b < s$. And as

$$-\frac{d-s}{d-s-g-b} = \mu(N) \geq \mu(M_{V,\mathcal{L}}) = -\frac{d}{d-g-c} ,$$

then $s(g+c) \leq d(c-b)$, hence $c-b > 0$ and $s \leq \frac{d}{g+c}(c-b)$.

2.1 Line bundles of degree $d = 2g + 2$

A first consequence of these lemmas is the following proposition asserting semistability for hyperplane tranforms of line bundles of degree $2g + 2$.

Proposition 2.5 *Let $\mathcal{L}$ be a line bundle of degree $d = 2g + 2$ on a curve C of genus $g \geq 2$. Then $M_{V,\mathcal{L}}$ is semistable for every generating hyperplane $V \subset H^0(\mathcal{L})$. It is strictly semistable if C is hyperelliptic.*

Proof Let us prove the semistability of $M_{V,\mathcal{L}}$.

Consider a stable subbundle $N \hookrightarrow M_{V,\mathcal{L}}$ of maximal slope, and suppose that it destabilizes $M_{V,\mathcal{L}}$ in the strict sense, *i.e.* $\mu(N) > -2 = \mu(M_{V,\mathcal{L}})$. By lemma 2.2 and remark 2.4 (we have $b = 0$ in this case), we know that N fits into a diagram

$$\begin{array}{ccccccccc} 0 & \rightarrow & N & \rightarrow & H^0(F) \otimes \mathcal{O}_C & \rightarrow & F & \rightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \rightarrow & M_{V,\mathcal{L}} & \rightarrow & V \otimes \mathcal{O}_C & \rightarrow & \mathcal{L} & \rightarrow & 0 \end{array}$$

with F a line bundle of degree $\deg F \leq d - 2 = 2g$. Moreover, $h^1(F) = 0$ since otherwise $\mu(N) \leq -2$ by lemma 2.1. Hence $\text{rk} N = \deg F - g$, and

$$\mu(N) = -\deg F / (\deg F - g) \leq -2g / (2g - g) = -2 ,$$

(again, use that the function $-x/(x-g)$ is strictly increasing for $x > g$). So it is not possible to find a strictly destabilizing N .

If the curve is hyperelliptic, then $M_{V,\mathcal{L}}$ is strictly semistable: we can show that there is a line bundle of degree -2 injecting in $M_{V,\mathcal{L}}$. In fact we can consider the line bundle A dual of the only g_2^1 of the curve, *i.e.* the dual of the hyperelliptic bundle.

The hyperelliptic bundle A^* has $h^0(A^*) = 2$, and from the exact sequence $0 \rightarrow M_{V,\mathcal{L}} \otimes A^* \rightarrow V \otimes A^* \rightarrow \mathcal{L} \otimes A^* \rightarrow 0$, we see that there are destabilizations of $M_{V,\mathcal{L}}$ by the line bundle A if and only if

$$H^0(M_{V,\mathcal{L}} \otimes A^*) = \ker(\varphi: V \otimes H^0(A^*) \rightarrow H^0(\mathcal{L} \otimes A^*)) \neq 0 .$$

Counting dimensions we see that the map φ cannot be injective:

$$\dim V \cdot \dim H^0(A^*) = (g+2)2 > g+5 = \dim H^0(\mathcal{L} \otimes A^*) .$$

□

In order to prove stability for non hyperelliptic curves though, we need to take a generic hyperplane, and not just a generating one.

The following is a special case of a more general result proven in section 2.3

Theorem 2.6 *Let $\mathcal{L}$ be a line bundle of degree $d = 2g + 2$ on a curve C of genus $g \geq 2$. Then $M_{V,\mathcal{L}}$ is stable for a generic hyperplane $V \subset H^0(\mathcal{L})$ if and only if C is non hyperelliptic.*

2.2 Line bundles of degree $d > 2g + 2c$

Here we show that for a generic subspace the transform of a line bundle of degree $d > 2g + 2c$ is stable. In contrast to proposition 2.5, we have to consider generic hyperplanes, and not just generating ones.

Theorem 2.7 *Let $\mathcal{L}$ be a line bundle of degree d on a curve C of genus $g \geq 2$, such that $d > 2g + 2c$, with $1 \leq c \leq g$. Then $M_{V,\mathcal{L}}$ is stable for a generic subspace $V \subset H^0(\mathcal{L})$ of codimension c .*

Proof

Let us proceed as in proposition 2.5. We have that $-2 < \mu(M_{V,\mathcal{L}}) < -1$.

Consider a stable subbundle $N \hookrightarrow M_{V,\mathcal{L}}$ of maximal slope. By lemma 2.2 we know it fits into a diagram

$$\begin{array}{ccccccc} 0 & \rightarrow & N & \rightarrow & W \otimes \mathcal{O}_C & \rightarrow & F \rightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \rightarrow & M_{V,\mathcal{L}} & \rightarrow & V \otimes \mathcal{O}_C & \rightarrow & \mathcal{L} \rightarrow 0. \end{array}$$

We can right away conclude that $h^1(F) = 0$, as by lemma 2.1 we would otherwise have $\mu(N) \leq -2$.

So F is a globally generated line bundle with $h^1(F) = 0$, $\deg F =: d - s \leq d - 2$, and W is a b -codimensional subspace of $H^0(F)$. By remark 2.4, we see that for every b with $0 \leq b < c$ there is a finite number of s giving rise to a possible destabilization of $M_{V,\mathcal{L}}$.

For any of those b and s we will construct a parameter space allowing F , W , and the subspace $V \subset H^0(\mathcal{L})$ to vary.

For any such b and s we want to consider the parameter space $\mathcal{D}_{b,s}$, parametrizing subspaces $V \subset H^0(\mathcal{L})$ together with a destabilizing bundle of $M_{V,\mathcal{L}}$ of degree $s - d$ originating from a subspace W as in the construction above:

$$\begin{aligned} \mathcal{D}_{b,s} := \{ & (F, F \hookrightarrow \mathcal{L}, W \subset H^0(F)), V \subset H^0(\mathcal{L}) \mid F \in \text{Pic}^{d-s}(C), \\ & (\varphi: F \hookrightarrow \mathcal{L}) \in \mathbb{P}(H^0(F^* \otimes \mathcal{L})), W \in \text{Gr}(b, H^0(F)) \\ & V \in \text{Gr}(c, H^0(\mathcal{L})), \varphi|_W: W \hookrightarrow V \subset H^0(\mathcal{L}) \} . \end{aligned}$$

In order to estimate its dimension, we use the natural morphisms

$$\pi_{b,s}: \mathcal{D}_{b,s} \rightarrow \text{Pic}^{d-s}(C), (F, F \hookrightarrow \mathcal{L}, W, V) \mapsto F,$$

and $\rho_{b,s}: \mathcal{D}_{b,s} \rightarrow \text{Gr}(c, H^0(\mathcal{L}))$, $(F, F \hookrightarrow \mathcal{L}, W, V) \mapsto V$.

The image of $\pi_{b,s}$ is formed by all the line bundles $F \in \text{Pic}^{d-s}(C)$ such that $h^0(F^* \otimes \mathcal{L}) \neq 0$. In particular $\dim \pi_s(\mathcal{D}_s) = \min(s, g)$, because the degree of $F^* \otimes \mathcal{L}$ is s . The fiber over $F \in \pi_{b,s}(\mathcal{D}_s)$ has the same dimension as $\mathbb{P}(H^0(F^* \otimes \mathcal{L})) \times \text{Gr}(b, (H^0(F))) \times \text{Gr}(c, (H^0(\mathcal{L})/W))$.

By Clifford's theorem, $h^0(F^* \otimes \mathcal{L}) = s/2 + 1$ if $s \leq 2g$, and $h^0(F^* \otimes \mathcal{L}) = s + 1 - g$ otherwise. So,

$$\dim \mathcal{D}_{b,s} \leq \min(s, g) + \sup(s/2, s - g) + b(d - s - g + 1 - b) + c(s + b - c) \leq \\ (3/2)s + b(d - s - g + 1 - b) + c(s + b - c) .$$

Claim: for g, d, c as in the hypothesis and s, b satisfying the inequalities of remark 2.4, we have

$$(3/2)s + b(d - s - g + 1 - b) + c(s + b - c) < c(d + 1 - g) - c^2 = \dim \text{Gr}(c, H^0(\mathcal{L})) .$$

Proving the claim, we show that for all s and b giving rise to possible destabilizations, the morphisms $\rho_{b,s}: \mathcal{D}_{b,s} \rightarrow \text{Gr}(c, H^0(\mathcal{L}))$ have a locally closed image of dimension strictly smaller than $\text{Gr}(c, H^0(\mathcal{L}))$, hence the generic subspace avoids all possible destabilizations of $M_{V, \mathcal{L}}$.

The claim is equivalent to

$$\frac{3s}{2(c-b)} + s + b < d + 1 - g ,$$

using inequalities (3) we get

$$\frac{3s}{2(c-b)} + s + b \leq \frac{3/2 + (c-b)}{g+c} d + b ,$$

hence we want to prove

$$\frac{3/2 + (c-b)}{g+c} d + b < d + 1 - g ,$$

which is equivalent to

$$\frac{b+g-1}{b+g-3/2} < \frac{d}{g+c} ,$$

and as $b \geq 0 \geq 2 - g$ then $\frac{b+g-1}{b+g-3/2} \leq 2 < \frac{d}{g+c}$. □

2.3 Line bundles of degree $d = 2g + 2c$

We have shown in section 2.1 that hyperplane transforms of a degree $2g + 2$ line bundle are always semistable.

We prove here, that generic c -codimensional transforms of a degree $2g + 2c$ line bundle are stable, except in the hyperelliptic case, where they are strictly semistable.

Theorem 2.8 *Let $\mathcal{L}$ be a line bundle of degree $d = 2g + 2c$ on a curve C of genus $g \geq 2$. Then $M_{V,\mathcal{L}}$ is semistable for a generic subspace $V \subset H^0(\mathcal{L})$ of codimension c . It is stable if and only if C is non hyperelliptic.*

Proof As in the proof of theorem 2.7 we want to construct parameter spaces for destabilizations, and verify by dimension count that the generic subspace avoids them.

Let us consider a line bundle $\mathcal{L}$ of degree $d = 2g + 2c$ on a curve C of genus $g \geq 2$, and the transform $M_{V,\mathcal{L}}$ for a subspace $V \subset H^0(\mathcal{L})$ of codimension c .

To show semistability, let us suppose that there is a destabilizing stable vector bundle $N \hookrightarrow M_{V,\mathcal{L}}$, with $\mu(N) > \mu(M_{V,\mathcal{L}}) = -2$.

By lemma 2.2 we know it fits in the diagram

$$\begin{array}{ccccccccc} 0 & \rightarrow & N & \rightarrow & W \otimes \mathcal{O}_C & \rightarrow & F & \rightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \rightarrow & M_{V,\mathcal{L}} & \rightarrow & V \otimes \mathcal{O}_C & \rightarrow & \mathcal{L} & \rightarrow & 0 . \end{array}$$

and we can suppose that $h^1(F) = 0$ by lemma 2.1.

In this case we can follow the same computations as in theorem 2.7: we have a parameter space for destabilizations

$$\begin{aligned} \mathcal{D}_{b,s} := \{ & (F, F \hookrightarrow \mathcal{L}, W \subset H^0(F)), V \subset H^0(\mathcal{L}) \mid F \in \text{Pic}^{d-s}(C), \\ & (\varphi: F \hookrightarrow \mathcal{L}) \in \mathbb{P}(H^0(F^* \otimes \mathcal{L})), W \in \text{Gr}(b, H^0(F)) \\ & V \in \text{Gr}(c, H^0(\mathcal{L})), \varphi|_W: W \hookrightarrow V \subset H^0(\mathcal{L}) \} , \end{aligned}$$

whose dimension is bounded by

$$\dim \mathcal{D}_{b,s} \leq (3/2)s + b(d - s - g + 1 - b) + c(s + b - c) ,$$

with b and s satisfying $0 < c - b < s \leq \frac{d}{g+c}(c - b)$.

Except in the case $b = 0$ and $g = 2$, we can follow the very same proof of theorem 2.7, and we see that this bound shows that the generic subspace avoids the destabilization locus.

In the case $b = 0$ and $g = 2$ as well, it can be easily shown that $\dim \mathcal{D}_{b,s} < \dim \text{Gr}(c, H^0(\mathcal{L}))$, for all s giving rise to destabilizations.

To show that we have strict semistability in the hyperelliptic case, we can proceed as in proposition 2.5, and show that dual of the hyperelliptic bundle is a subbundle of $M_{V,\mathcal{L}}$, of slope -2 .

To show that we have stability in the non hyperelliptic case, we have to exclude slope -2 subbundles $N \hookrightarrow M_{V,\mathcal{L}}$.

Again we can apply lemma 2.2 and consider the diagram

$$\begin{array}{ccccccccc} 0 & \rightarrow & N & \rightarrow & W \otimes \mathcal{O}_C & \rightarrow & F & \rightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \rightarrow & M_{V,\mathcal{L}} & \rightarrow & V \otimes \mathcal{O}_C & \rightarrow & \mathcal{L} & \rightarrow & 0 , \end{array}$$

where we can distinguish the two cases $H^1(F) = 0$, and $H^1(F) \neq 0$.

In the case $H^1(F) = 0$ we can follow again the same computations as in theorem 2.7.

In the case $H^1(F) \neq 0$, lemma 2.1 implies $F = \omega$ and $N = M_\omega$, hence the parameter space for destabilizations will be

$$\mathcal{D} := \{(\omega \hookrightarrow \mathcal{L} , V \subset H^0(\mathcal{L})) \mid H^0(\omega) \subset V\} ,$$

and it can be shown that $\dim \mathcal{D} < \dim \mathrm{Gr}(c, H^0(\mathcal{L}))$.

□

3 Theta divisors and transforms

When a vector bundle has integer slope $\mu(E) = \mu \in \mathbb{Z}$, we can define the set

$$\Theta_E := \{P \in \mathrm{Pic}^\nu(C) \mid H^0(C, E \otimes P) \neq 0\} ,$$

where $\nu := g - 1 - \mu$.

As $\chi(E \otimes P) = 0$, either $\Theta_E = \mathrm{Pic}^\nu(C)$, or it has a natural structure of effective divisor in $\mathrm{Pic}^\nu(C)$. In the latter case we say that E *admits a theta divisor*. The class of this divisor in $H^2(\mathrm{Pic}^\nu(C), \mathbb{Z})$ is $\mathrm{rk} E \cdot \vartheta$, where ϑ is the class of the canonical theta divisor of $\mathrm{Pic}^\nu(C)$.

Whenever a vector bundle admits a theta divisor, then it is semistable. And strictly semistable vector bundles admitting a theta divisor have *non integral* theta divisors.

However, there are examples of stable vector bundles with no theta divisor, or with a reducible theta divisor.

Beauville shows in [Bea03] that the total transform M_L of a degree $2g$ line bundle L on a genus g curve C always has a reducible theta divisor. And that if L is very ample, and C is not hyperelliptic, then M_L is stable.

The vector bundles considered above, *i.e.* transforms of degree $d \geq 2g + 2c$ line bundles, with respect to c codimensional subspaces of global sections, have slope μ such that $-2 \leq \mu < -1$. The case of integer slope $\mu = -2$ appears if and only if $d = 2g + 2c$.

Following the same argument as in [Bea03], we prove that for the generic $V \subset H^0(C, \mathcal{L})$ those transforms always carry a non integral theta divisor.

To prove that, for a generic $V \subset H^0(C, \mathcal{L})$ within the numerical conditions above, the transform $M_{V, \mathcal{L}}$ admits a theta divisor, we need the following lemma:

Lemma 3.1 *Let P be a 2-dimensional vector space, H a vector space of dimension $n + c$, and $K \subset P \otimes H$ a subspace of dimension $2c$. If K contains no pure vectors, then the generic n -dimensional subspace $V \in \mathrm{Gr}(n, H)$ verifies*

$$K \cap (P \otimes V) = 0 .$$

Proof We consider the map

$$\begin{array}{ccc} f: \mathrm{Gr}(n, H) & \rightarrow & \mathrm{Gr}(2n, P \otimes H) \\ V & \mapsto & P \otimes V , \end{array}$$

and we claim that the image of f is not contained in the closed subscheme

$$Z := \{W \in \text{Gr}(2n, P \otimes H) \mid \dim K \cap W \geq 1\} .$$

Let us observe at first that Z carries a filtration

$$Z = Z_1 \supseteq Z_2 \supseteq \dots \supseteq Z_s := \{W \subset P \otimes H \mid \dim K \cap W \geq s\} \supseteq \dots .$$

The tangent space of the grassmannian $\text{Gr}(2n, P \otimes H)$ at a point W is

$$T_W \text{Gr}(2n, P \otimes H) = \text{Hom}(W, P \otimes H/W) .$$

The subscheme $Z_s \setminus Z_{s+1}$ is smooth and its tangent space at a point W , is given by first order deformations of $W \subset P \otimes H$ that deform $W \cap K$ into an s -dimensional subspace of K :

$$T_W(Z_s \setminus Z_{s+1}) = \{\varphi \in \text{Hom}(W, P \otimes H/W) \mid \varphi(W \cap K) \subseteq K/(W \cap K)\} .$$

And the differential of the morphism f at the point $V \in \text{Gr}(n, H)$ is the map

$$\begin{aligned} df_V : \quad T_V \text{Gr}(n, H) &\rightarrow T_{P \otimes V} \text{Gr}(2n, P \otimes H) \\ \varphi \in \text{Hom}(V, H/V) &\mapsto 1 \otimes \varphi \in \text{Hom}(P \otimes V, P \otimes (H/V)) . \end{aligned}$$

We can prove now that if $V \in \text{Gr}(n, H)$ is a subspace such that $P \otimes V \in Z_s \setminus Z_{s+1}$, then $df_V(T_V \text{Gr}(n, H)) \not\subseteq T_{P \otimes V}(Z_s \setminus Z_{s+1})$: we claim that there exists a $\varphi \in \text{Hom}(V, H/V)$ such that $1 \otimes \varphi \notin T_{P \otimes V}(Z_s \setminus Z_{s+1})$.

To see this, let us choose a basis (e_1, e_2) for P , and a vector $w = e_1 \otimes v_1 + e_2 \otimes v_2 \in K \cap (P \otimes V)$. By the hypothesis on K , $v_1 \nparallel v_2$. Let us consider now a vector $z = e_1 \otimes z_1 + e_2 \otimes z_2 \in (P \otimes (H/V))$ such that $z \notin (K/(P \otimes V \cap K))$.

Then if we choose a $\varphi \in \text{Hom}(V, H/V)$ such that $\varphi(v_1) = z_1$ and $\varphi(v_2) = z_2$, we have that $(1 \otimes \varphi)(w) = z \notin K/(P \otimes V \cap K)$. Hence the image of a generic deformation of V avoids the subscheme $Z \subset \text{Gr}(2n, P \otimes H)$. $\square$

We can now prove the existence of theta divisors for generic transforms of slope -2 .

Theorem 3.2 *Let $\mathcal{L}$ be a line bundle of degree $d = 2g + 2c$ on a genus g curve C , where $c \in \mathbb{N}$ is a positive integer and $g \geq 2$. Then, if $V \subset H^0(C, \mathcal{L})$ is a generic c -codimensional subspace, the transform $M_{V, \mathcal{L}}$ admits a non integral theta divisor.*

Proof We recall that $\mu(M_{V, \mathcal{L}}) = -2$. We have to show first that, for the generic $V \subset H^0(C, \mathcal{L})$, $\Theta_{M_{V, \mathcal{L}}} \neq \text{Pic}^{g+1}(C)$, i.e. that there is a $P \in \text{Pic}^{g+1}(C)$ such that $H^0(M_{V, \mathcal{L}} \otimes P) = 0$. By the exact sequence

$$0 \rightarrow M_{V, \mathcal{L}} \otimes P \rightarrow V \otimes P \rightarrow \mathcal{L} \otimes P \rightarrow 0$$

this is the same as a $P \in \text{Pic}^{g+1}(C)$ such that the multiplication map

$$\mu : V \otimes H^0(P) \rightarrow H^0(\mathcal{L} \otimes P)$$

is injective.

If P belongs to the divisor $D = (\omega_C) - C_{g-2} + C \subset \text{Pic}^{g+1}(C)$, i.e. if P can be written in the form $P = \omega_C(x_1 - x_2 - \dots - x_{g-1})$ for some points $x_1, x_2, \dots, x_{g-1} \in C$, then either $h^0(P) > 2$, or $h^0(P) = 2$ and P has a base point. In any case this implies that μ is not injective for any V (c.f. [Bea03]).

Any P in $\text{Pic}^{g+1}(C) \setminus D$ is base point free and has $h^0(P) = 2$. Let us fix such a P , and assume by generality that $h^1(L \otimes P^*) = 0$. We claim that for the generic $V \subset H^0(C, \mathcal{L})$ of codimension c the multiplication map $\mu: H^0(P) \otimes V \rightarrow H^0(P \otimes \mathcal{L})$ is injective. From the exact sequence

$$0 \rightarrow P^* \rightarrow H^0(P) \otimes \mathcal{O}_C \rightarrow P \rightarrow 0$$

we get

$$0 \rightarrow H^0(P^* \otimes \mathcal{L}) \rightarrow H^0(P) \otimes H^0(\mathcal{L}) \rightarrow H^0(P \otimes \mathcal{L}) \rightarrow 0,$$

hence the map μ is injective if and only if the subspace $V \subset H^0(C, \mathcal{L})$ verifies $H^0(P^* \otimes \mathcal{L}) \cap (H^0(P) \otimes V) = 0$. And this is given by lemma 3.1.

Hence we know that for the generic subspace V , the transform $M_{V, \mathcal{L}}$ admits a theta divisor. To observe that it is not integral, we notice that the set of points of $\Theta_{M_{V, \mathcal{L}}}$ contains the divisor D whose cohomology class is $(g-1)\vartheta$ (cf. [FMP03]). As the cohomology class of $\Theta_{M_{V, \mathcal{L}}}$ is $(g+c)\vartheta$, it must be a non integral divisor. □

As we have proved the existence of theta divisors for transforms with respect to subspaces of any codimension, this shows semistability in some cases not previously treated:

Corollary 3.3 *Let $\mathcal{L}$ be a line bundle of degree $d = 2g + 2c$ on a genus g curve C , where $c \in \mathbb{N}$ is any positive integer and $g \geq 2$. Then, if $V \subset H^0(C, \mathcal{L})$ is a generic c -codimensional subspace, the transform $M_{V, \mathcal{L}}$ is semistable.*

Remark 3.4 If C is not hyperelliptic and $\mathcal{L}$ is a degree $d = 2g + 2c$ line bundle, where $d \notin 2(g-1)\mathbb{N}$, then the transform $M_{V, \mathcal{L}}$ of $\mathcal{L}$ with respect to a generic subspace $V \subset H^0(C, \mathcal{L})$ of codimension c admits a reducible theta divisor. In fact the set of points of $\Theta_{M_{V, \mathcal{L}}}$ contains the divisor

$$D = (\omega_C) - C_{g-2} + C \subset \text{Pic}^{g+1}(C),$$

which is irreducible if C is not hyperelliptic, and whose cohomology class is $(g-1)\vartheta$. As the cohomology class of $\Theta_{M_{V, \mathcal{L}}}$ is $(g+c)\vartheta = (d/2)\vartheta$, it cannot be a multiple of $(g-1)\vartheta$, then $\Theta_{M_{V, \mathcal{L}}}$ must be reducible.

Hence, if $c \leq g$ and $c \neq g-2$ we have further examples of stable vector bundles (by theorem 2.8) with reducible theta divisors.

4 Conclusions

We have proven stability of transforms of line bundles with respect to subspaces of low codimension. On the converse, it is rather easy to show the stability of transforms with respect to subspaces of low dimension: any stable vector bundle M^* of slope $\mu(M^*) > 2g - 1$ is globally generated. Hence we can pick any stable vector bundle M^* of determinant $\mathcal{L}$ and rank r , such that $r < d/(2g - 1)$, where $\deg \mathcal{L} = d$. Choosing any generating subspace $V^* \subset H^0(M^*)$ of rank $r + 1$, we get an exact sequence

$$0 \rightarrow \mathcal{L}^* \rightarrow V^* \otimes \mathcal{O} \rightarrow M^* \rightarrow 0 .$$

Dualizing we get an exact sequence

$$0 \rightarrow M \rightarrow V \otimes \mathcal{O} \rightarrow \mathcal{L} \rightarrow 0 ,$$

where M is a stable transform of $\mathcal{L}$. Hence, every stable bundle of rank $r < d/(2g - 1)$ and determinant $\mathcal{L}^*$, is a stable transform of $\mathcal{L}$. So the rational map $\mathbb{G}r(r + 1, H^0(\mathcal{L})) \dashrightarrow SU(r, \mathcal{L})$ is dominant.

By the same argument we see that there is only one globally generated vector bundle, among vector bundles of determinant $\mathcal{L}$ and rank $d - g$ with no trivial summands, where $d = \deg \mathcal{L} \geq 2g$. Furthermore this is semistable, and even stable if $d > 2g$. In fact having such a globally generated bundle N , we can pick a vector space V of global sections of dimension $\text{rk} N + 1$ generating N . This gives rise to the exact sequence

$$0 \rightarrow \mathcal{L}^* \rightarrow V \otimes \mathcal{O} \rightarrow N \rightarrow 0 ,$$

and dualizing

$$0 \rightarrow N^* \rightarrow V^* \otimes \mathcal{O} \rightarrow \mathcal{L} \rightarrow 0 .$$

But as N is globally generated and has no trivial summands, then $H^0(N^*) = 0$. And since V^* and $H^0(\mathcal{L})$ have the same dimension, then $V^* \xrightarrow{\sim} H^0(\mathcal{L})$. Hence $N^* = M_{\mathcal{L}}$ is unique.

So when we consider the rational map $\mathbb{G}r(r + 1, H^0(\mathcal{L})) \dashrightarrow SU(r, \mathcal{L})$, $V \mapsto (M_{V, \mathcal{L}})^*$, we are saying that its image is made by globally generated bundles, and we can sum all this up in the following table, where we suppose that $d > 2g + 2c$, with $1 \leq c \leq g$:

$\text{rk}(M_{V, \mathcal{L}}) = r$	stability	map
$1 \leq r < d/(2g - 1)$	stable	$\mathbb{G}r(r + 1, H^0(\mathcal{L})) \dashrightarrow SU(r, \mathcal{L})$ dominant
$\frac{d}{2g-1} \leq r < d - g - c$	??	??
$d - g - c \leq r < d - g$	stable	$\mathbb{G}r(r + 1, H^0(\mathcal{L})) \dashrightarrow SU(r, \mathcal{L})$
$r = d - g$	stable	$\{*\} \hookrightarrow SU(r, \mathcal{L})$

where theorem 2.7 corresponds to the existence of the rational map

$$\mathbb{G}r(r + 1, H^0(\mathcal{L})) \dashrightarrow SU(r, \mathcal{L}) .$$

References

- [Bea03] Arnaud Beauville, *Some stable vector bundles with reducible theta divisor*, Manuscripta Math. **110** (2003), no. 3, 343–349.
- [But94] David C. Butler, *Normal generation of vector bundles over a curve*, J. Differential Geom. **39** (1994), no. 1, 1–34.
- [EL92] Lawrence Ein and Robert Lazarsfeld, *Stability and restrictions of Picard bundles, with an application to the normal bundles of elliptic curves*, Complex projective geometry (Trieste, 1989/Bergen, 1989), London Math. Soc. Lecture Note Ser., vol. 179, Cambridge Univ. Press, Cambridge, 1992, pp. 149–156.
- [FMP03] Gavril Farkas, Mircea Mustață, and Mihnea Popa, *Divisors on $\mathcal{M}_{g,g+1}$ and the minimal resolution conjecture for points on canonical curves*, Ann. Sci. École Norm. Sup. (4) **36** (2003), no. 4, 553–581.
- [Mer99] Vincent Mercat, *Le problème de Brill-Noether pour des fibrés stables de petite pente*, J. Reine Angew. Math. **506** (1999), 1–41.
- [PR88] Kapil Paranjape and S. Ramanan, *On the canonical ring of a curve*, Algebraic geometry and commutative algebra, Vol. II, Kinokuniya, Tokyo, 1988, pp. 503–516.

Ernesto Carlo MISTRETTA
ernesto@math.jussieu.fr
INSTITUT DE MATHÉMATIQUES DE JUSSIEU
Équipe de Topologie et Géométrie Algébriques
175, rue de Chevaleret
75013 Paris